

IGUSA'S p -ADIC LOCAL ZETA FUNCTION ASSOCIATED TO A POLYNOMIAL MAPPING AND A POLYNOMIAL INTEGRATION MEASURE

BART BORIES

ABSTRACT. For p prime, we give an explicit formula for Igusa's local zeta function associated to a polynomial mapping $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$, with $f_1, \dots, f_t \in \mathbf{Z}_p[x_1, \dots, x_n]$, and an integration measure on \mathbf{Z}_p^n of the form $|g(x)||dx|$, with g another polynomial in $\mathbf{Z}_p[x_1, \dots, x_n]$. We treat the special cases of a single polynomial and a monomial ideal separately. The formula is in terms of Newton polyhedra and will be valid for \mathbf{f} and g sufficiently non-degenerated over \mathbf{F}_p with respect to their Newton polyhedra. The formula is based on, and is a generalization of results in [6], [9] and [15].

CONTENTS

Introduction	1
Acknowledgements	4
1. Preliminaries	4
2. Igusa's zeta function of a monomial ideal and a polynomial measure	7
3. Computation of an important integral using Hensel's Lemma	12
4. Igusa's zeta function of a single polynomial and a polynomial measure	16
5. Igusa's zeta function of a polynomial mapping and a polynomial measure	18
References	20

INTRODUCTION

Motivation. In order to prove the rationality of a generating series counting the number of solutions of polynomial congruences, in 1974 Igusa [10] introduced his p -adic zeta function. In 1992, Denef and Loeser [7] obtained the topological zeta function as a kind of limit of p -adic zeta functions; later they presented in [8] the finer and intrinsically defined motivic zeta function. All these invariants were originally associated to one polynomial or analytic function in several variables over a certain field (a p -adic field, \mathbf{C} and an arbitrary field of characteristic zero, respectively), but the concepts have been generalized in many ways during the research that followed. For instance, we now associate these zeta functions to several polynomials or to an ideal in a polynomial ring, see e.g. [9, 15, 13, 14]. Nowadays one is also interested in zeta functions of a number of polynomials and a more general measure or differential form. For example, in [11] Veys and Némethi use this notion

Date: February 28, 2012.

to prove a remarkable result on the generalized monodromy conjecture. For other applications, see e.g. [1, 2]. In this respect, it is useful to examine which known results and formulas have their analogue in this generalized context. In this paper, we adapt results from [6, 9, 15] and prove an explicit formula for Igusa's p -adic zeta function of a strongly non-degenerated polynomial mapping and a 'polynomial measure'. Although we state the formula for the p -adic one, the result can also be formulated for the topological and motivic zeta functions. In [3] we will use this formula to investigate the analogue of Veys and Némethi's result for a related, stronger conjecture of Igusa–Denef–Loeser.

Igusa's zeta function of a polynomial mapping and a polynomial measure.

For a prime p , we denote by \mathbf{Q}_p the field of p -adic numbers and by \mathbf{Z}_p its subring of p -adic integers. Denote by $|\cdot|$ the p -adic norm on \mathbf{Q}_p . On \mathbf{Q}_p^n , $n \in \mathbf{N} \setminus \{0\}$, we consider the Haar measure, so normalized that \mathbf{Z}_p^n has measure one, and we denote it by $|dx| = |dx_1 \wedge \cdots \wedge dx_n|$. For a measurable subset $A \subset \mathbf{Q}_p^n$, we denote its measure by $\mu(A)$. Let us start with the definition of Igusa's local zeta function.

Definition 0.1 (Igusa's p -adic local zeta function). Let p be a prime number and $f(x) = f(x_1, \dots, x_n)$ a polynomial in $\mathbf{Q}_p[x_1, \dots, x_n]$. Igusa's (p -adic) local zeta function associated to f is defined as

$$Z_f : \{s \in \mathbf{C} \mid \Re(s) > 0\} \rightarrow \mathbf{C} : s \mapsto Z_f(s) = \int_{\mathbf{Z}_p^n} |f(x)|^s |dx|.$$

More generally, one defines Igusa's local zeta function associated to a polynomial mapping $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$, with $f_1, \dots, f_t \in \mathbf{Q}_p[x_1, \dots, x_n]$, as

$$(1) \quad Z_{\mathbf{f}} : \{s \in \mathbf{C} \mid \Re(s) > 0\} \rightarrow \mathbf{C} : s \mapsto Z_{\mathbf{f}}(s) = \int_{\mathbf{Z}_p^n} \|\mathbf{f}(x)\|^s |dx|,$$

where $\|\mathbf{f}(x)\| = \max_i |f_i(x)|$.

To an ideal \mathcal{I} of $\mathbf{Z}_p[x_1, \dots, x_n]$, we associate Igusa's local zeta function

$$Z_{\mathcal{I}} : \{s \in \mathbf{C} \mid \Re(s) > 0\} \rightarrow \mathbf{C} : s \mapsto Z_{\mathcal{I}}(s) = \int_{\mathbf{Z}_p^n} |\mathcal{I}(x)|^s |dx|,$$

where $|\mathcal{I}(x)| = \max\{|f(x)| \mid f \in \mathcal{I}\}$.

We can further generalize these notions by considering alternative, 'polynomial' integration measures on \mathbf{Z}_p^n , i.e. measures of the form $|g(x)||dx|$, with g another polynomial in $\mathbf{Q}_p[x_1, \dots, x_n]$. For f , \mathbf{f} and \mathcal{I} as above, we put

$$Z_{f,gdx}(s) = \int_{\mathbf{Z}_p^n} |f(x)|^s |g(x)||dx|, \quad Z_{\mathbf{f},gdx}(s) = \int_{\mathbf{Z}_p^n} \|\mathbf{f}(x)\|^s |g(x)||dx|$$

and $Z_{\mathcal{I},gdx}(s) = \int_{\mathbf{Z}_p^n} |\mathcal{I}(x)|^s |g(x)||dx|.$

These are the zeta functions that we will study here.

Remark 0.2. In general, we associate p -adic zeta functions to polynomial mappings $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$, rather than to ideals $(f_1, \dots, f_t) \triangleleft \mathbf{Q}_p[x_1, \dots, x_n]$, since $Z_{\mathbf{f}}$, as defined in (1), depends on the specific polynomials f_1, \dots, f_t , not only on the ideal generated by them. However, if we consider polynomials f_1, \dots, f_t over the p -adic integers, it turns out that $Z_{\mathbf{f}}$ only depends on the ideal $(f_1, \dots, f_t) \triangleleft \mathbf{Z}_p[x_1, \dots, x_n]$, i.e. two polynomial mappings over \mathbf{Z}_p give rise to the same zeta

function, if their composing polynomials generate the same ideal of $\mathbf{Z}_p[x_1, \dots, x_n]$. This yields a proper definition of Igusa's zeta function of a $\mathbf{Z}_p[x_1, \dots, x_n]$ ideal, which coincides with the one given above. Indeed, for every set $\{f_1, \dots, f_t\}$ of generators of an ideal $\mathcal{I} \triangleleft \mathbf{Z}_p[x_1, \dots, x_n]$, we have that

$$|\mathcal{I}(x)| = \max_{f \in \mathcal{I}} |f(x)| = \max_{1 \leq i \leq t} |f_i(x)|$$

for every $x \in \mathbf{Z}_p^n$.

Following [9], we will use the terminology of ideals when it comes to monomial ideals (Sections 1–2), otherwise we will speak in terms of polynomial mappings.

Using resolution of singularities, Igusa [10] proved in 1974 that his p -adic zeta function Z_f , associated to one polynomial f , is a rational function in the variable $t = p^{-s}$. An alternative proof (based on p -adic cell decomposition) of this important fact was obtained ten years later by Denef [5]. With the same techniques it can be proved that this rationality result still holds for the other versions of Igusa's local zeta function, described above. Hence, all of these functions allow a meromorphic continuation to the whole complex plane, that we denote with the same symbol.

Considering Newton polyhedra (see Definition 1.2 below), Denef and Hoornaert obtained in their 2001 paper [6] a very explicit formula for Igusa's local zeta function of a single polynomial, for a special class of polynomials, namely those who are non-degenerated over \mathbf{F}_p with respect to all the faces of their Newton polyhedron (see Definition 1.4). In their paper [9] of 2007, Howald, Mustață and Yuen prove a similar formula for Igusa's local zeta function $Z_{\mathcal{I}}$, associated to a monomial ideal $\mathcal{I} \triangleleft \mathbf{Z}[x_1, \dots, x_n]$ and the usual integration measure. In 2008, Veys and Zúñiga-Galindo [15] generalized those two formulas to a formula for Igusa's local zeta function $Z_{\mathbf{f}}$ of a polynomial mapping \mathbf{f} , that is strongly non-degenerated over \mathbf{F}_p with respect to its Newton polyhedron (Definition 1.5).

Our goal is to adapt those formulas to cover the generalized versions $Z_{f,gdx}$, $Z_{\mathbf{f},gdx}$ and $Z_{\mathcal{I},gdx}$ of Igusa's local zeta function, associated to a polynomial measure $|g(x)||dx|$ on \mathbf{Z}_p^n , where g is a polynomial in $\mathbf{Z}_p[x_1, \dots, x_n]$ that is also non-degenerated over \mathbf{F}_p with respect to its Newton polyhedron, and where the pairs (f, g) and (\mathbf{f}, g) satisfy a supplementary non-degeneracy condition, described in Definitions 4.2 and 5.5, respectively.

Overview. In the first section, we list all the needed definitions, notations and results that are already known, without much explanation. For more details on Igusa's zeta function and Newton polyhedra, we refer to [6], [9] and [15], for more background on convex geometry, we refer to [12]. In Section 2, we derive a formula for Igusa's zeta function associated to a monomial ideal and a measure $|g(x)||dx|$. A formula for Igusa's local zeta function of a single polynomial f and a measure $|g(x)||dx|$, is proved in Section 4. Section 3 is preparation for this section, here we derive a formula for the same integral as appearing in the definition of $Z_{f,gdx}(s)$, but with integration domain $(\mathbf{Z}_p^\times)^n$ instead of \mathbf{Z}_p^n . The proof of this formula is based on Hensel's Lemma. In the last section, we state the most general formula, i.e. for Igusa's zeta function of a polynomial mapping and a polynomial integration measure.

All definitions and results mentioned in this paper have straightforward analogues over any p -adic field, i.e. any field K , such that $[K : \mathbf{Q}_p] < \infty$.

ACKNOWLEDGEMENTS

The author would like to thank Wim Veys for proposing the problem and for many useful suggestions. He also wants to thank the referee for reviewing the paper and for some interesting comments.

1. PRELIMINARIES

Remark 1.1. In what follows, we will consider polynomials over \mathbf{Z}_p , because this is more convenient when dealing with the non-degeneracy conditions we use throughout the paper. Note that this does not affect the generality, since every polynomial f over \mathbf{Q}_p can be written as $f = p^{-i} \tilde{f}$ for some $i \in \mathbf{Z}_{\geq 0}$ and some $\tilde{f} \in \mathbf{Z}_p[x_1, \dots, x_n]$, leading to e.g. $Z_f(s) = p^{is} Z_{\tilde{f}}(s)$.

Definition 1.2 (Newton polyhedron). Let p be a prime number. We denote, for $\omega = (\omega_1, \dots, \omega_n) \in \mathbf{N}^n$, by x^ω the corresponding monomial $x_1^{\omega_1} \cdots x_n^{\omega_n}$ in $\mathbf{Z}_p[x_1, \dots, x_n]$. Let $f(x) = f(x_1, \dots, x_n) = \sum_{\omega \in \mathbf{N}^n} a_\omega x^\omega$ be a nonzero polynomial over \mathbf{Z}_p , satisfying $f(0) = 0$. Let $\mathbf{R}_{\geq 0} = \{x \in \mathbf{R} \mid x \geq 0\}$ and $\text{supp}(f) = \{\omega \in \mathbf{N}^n \mid a_\omega \neq 0\}$, the support of f . The Newton polyhedron Γ_f of f is defined as the convex hull in $\mathbf{R}_{\geq 0}^n$ of the set

$$\bigcup_{\omega \in \text{supp}(f)} \omega + \mathbf{R}_{\geq 0}^n.$$

The global Newton polyhedron Γ_f^{gl} of f is defined as the convex hull of $\text{supp}(f)$. Clearly, we have $\Gamma_f = \Gamma_f^{\text{gl}} + \mathbf{R}_{\geq 0}^n$.

Let $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$ be a nonconstant polynomial mapping, with $f_1, \dots, f_t \in \mathbf{Z}_p[x_1, \dots, x_n]$, satisfying $\mathbf{f}(0) = 0$. We define the support $\text{supp}(\mathbf{f})$ of \mathbf{f} as the union of the supports of its composing polynomials, and its Newton polyhedron $\Gamma_{\mathbf{f}}$, as above, as the convex hull in $\mathbf{R}_{\geq 0}^n$ of $\bigcup_{\omega \in \text{supp}(\mathbf{f})} \omega + \mathbf{R}_{\geq 0}^n$.

Let $\mathcal{I} \triangleleft \mathbf{Z}_p[x_1, \dots, x_n]$ be a nonzero proper monomial ideal¹. We define the Newton polyhedron $\Gamma_{\mathcal{I}}$ of \mathcal{I} as the convex hull in $\mathbf{R}_{\geq 0}^n$ of those $\omega \in \mathbf{N}^n$, such that $x^\omega \in \mathcal{I}$. The polyhedron $\Gamma_{\mathcal{I}}$ coincides with the Newton polyhedron of the monomial mapping $(x^{\omega_1}, \dots, x^{\omega_t})$ for any set $\{x^{\omega_1}, \dots, x^{\omega_t}\}$ of monomial generators of \mathcal{I} .

Notation 1.3. For $a \in \mathbf{Z}_p^n$, we denote by $\bar{a} = a + (p\mathbf{Z}_p)^n \in \mathbf{F}_p^n$ its reduction modulo $(p\mathbf{Z}_p)^n$. For $f \in \mathbf{Z}_p[x_1, \dots, x_n]$, we denote by \bar{f} the polynomial over \mathbf{F}_p , obtained from f , by reducing each of its coefficients modulo $p\mathbf{Z}_p$. Analogously, for a polynomial mapping $\mathbf{f} = (f_1, \dots, f_t)$, we denote $\bar{\mathbf{f}} = (\bar{f}_1, \dots, \bar{f}_t)$.

Definition 1.4 (Non-degenerated). Let f be as in Definition 1.2. For every face² τ of the Newton polyhedron Γ_f of f , we put

$$f_\tau(x) = \sum_{\omega \in \tau} a_\omega x^\omega.$$

We say that f is non-degenerated over \mathbf{F}_p with respect to (all the faces of) its Newton polyhedron Γ_f , if for every³ face τ of Γ_f , the locus of the polynomial \bar{f}_τ

¹A monomial ideal is an ideal that can be generated by monic monomials.

²By a face of Γ_f we mean Γ_f itself or one of its proper faces, which are the intersections of Γ_f with a supporting hyperplane (see e.g. [12]).

³Thus also for Γ_f .

has no singularities in $(\mathbf{F}_p^\times)^n$, or, equivalently, the set of congruences

$$\begin{cases} f_\tau(x) \equiv 0 \pmod{p} \\ \frac{\partial f_\tau}{\partial x_i}(x) \equiv 0 \pmod{p}, \quad i = 1, \dots, n \end{cases}$$

has no solutions in $(\mathbf{Z}_p^\times)^n$.

Definition 1.5 (Strongly non-degenerated). Let \mathbf{f} be as in Definition 1.2. For every face τ of the Newton polyhedron $\Gamma_{\mathbf{f}}$ of \mathbf{f} , we denote $\mathbf{f}_\tau = (f_{1,\tau}, \dots, f_{t,\tau})$, with the $f_{i,\tau}$ defined in the same way as the f_τ in Definition 1.4.

We call the mapping \mathbf{f} strongly non-degenerated over the field \mathbf{F}_p with respect to (all the faces of) its Newton polyhedron $\Gamma_{\mathbf{f}}$, if for every face τ of $\Gamma_{\mathbf{f}}$ and all $a \in (\mathbf{Z}_p^\times)^n$, satisfying $\mathbf{f}_\tau(a) \equiv 0 \pmod{p}$, the Jacobian matrix $J(\mathbf{f}_\tau, a) = ((\partial f_{i,\tau}/\partial x_j)(a))_{i,j}$ has maximal rank (equal to $\min(t, n)$) modulo p .⁴ In other words: \mathbf{f} is strongly non-degenerated over \mathbf{F}_p with respect to $\Gamma_{\mathbf{f}}$, if for every face τ of $\Gamma_{\mathbf{f}}$ and all $\bar{a} \in (\mathbf{F}_p^\times)^n$ in the locus of $\bar{\mathbf{f}}_\tau$, the Jacobian matrix $J(\bar{\mathbf{f}}_\tau, \bar{a}) = ((\partial \bar{f}_{i,\tau}/\partial x_j)(\bar{a}))_{i,j}$ over \mathbf{F}_p , has maximal rank.

Definition 1.6 ($m_f(k)$). Let f be as in Definition 1.2. For $k \in \mathbf{R}_{\geq 0}^n$, we define

$$m_f(k) = \inf_{x \in \Gamma_f} k \cdot x,$$

where $k \cdot x$ denotes the scalar product of k and x .

The infimum in the definition above is actually a minimum, where the minimum can as well be taken over the global Newton polyhedron Γ_f^{gl} of f , which is a compact set, or even over the finite set $\text{supp}(f)$.

Definition 1.7 (First meet locus). Let f be as in Definition 1.2 and $k \in \mathbf{R}_{\geq 0}^n$. We define the first meet locus of k as the face

$$F_f(k) = \{x \in \Gamma_f \mid k \cdot x = m_f(k)\}$$

of Γ_f .

Definition 1.8 (Primitive vector). A vector $k \in \mathbf{R}^n$ is called primitive if the components of k are integers whose greatest common divisor is one.

Definition 1.9 (Δ_f). Let f be as in Definition 1.2. For a face τ of Γ_f , we call

$$\Delta_f(\tau) = \{k \in \mathbf{R}_{\geq 0}^n \mid F_f(k) = \tau\}$$

the cone associated to τ . The $\Delta_f(\tau)$ are the equivalence classes of the equivalence relation \sim_f on $\mathbf{R}_{\geq 0}^n$, defined by

$$k \sim_f k' \quad \text{if and only if} \quad F_f(k) = F_f(k').$$

The ‘cones’ $\Delta_f(\tau)$ thus form a partition of $\mathbf{R}_{\geq 0}^n$ which we denote by Δ_f , so

$$\Delta_f = \{\Delta_f(\tau) \mid \tau \text{ is a face of } \Gamma_f\} = \mathbf{R}_{\geq 0}^n / \sim_f.$$

⁴Meaning that the matrix $((\partial f_{i,\tau}/\partial x_j)(a))_{i,j}$ over \mathbf{F}_p has maximal rank.

The $\Delta_f(\tau)$ are in fact relatively open^{5,6} convex cones⁷ with a very specific structure, as stated in the following lemma.

Lemma 1.10. [6, Lemma 2.6]. *Let f be as in Definition 1.2. Let τ be a proper face of Γ_f and let $\gamma_1, \dots, \gamma_r$ be the facets⁸ of Γ_f that contain τ . Let k_1, \dots, k_r be the unique primitive vectors in $\mathbf{N}^n \setminus \{0\}$ that are perpendicular to $\gamma_1, \dots, \gamma_r$, respectively. Then the cone $\Delta_f(\tau)$ associated to τ is the convex cone*

$$\Delta_f(\tau) = \{\lambda_1 k_1 + \lambda_2 k_2 + \dots + \lambda_r k_r \mid \lambda_i \in \mathbf{R}_{>0}\}$$

and its dimension⁹ equals $n - \dim \tau$.

Definition 1.11. For $k_1, \dots, k_r \in \mathbf{R}^n \setminus \{0\}$, we call $\Delta = \{\lambda_1 k_1 + \lambda_2 k_2 + \dots + \lambda_r k_r \mid \lambda_i \in \mathbf{R}_{>0}\}$ the cone strictly positively spanned by the vectors k_1, \dots, k_r . When the k_1, \dots, k_r can be chosen from \mathbf{Z}^n , we call it a rational cone. If we can choose k_1, \dots, k_r linearly independent over \mathbf{R} , Δ is called a simplicial cone. If Δ is rational and k_1, \dots, k_r can be chosen from a \mathbf{Z} -module basis of \mathbf{Z}^n , we call Δ a simple cone.

It follows from Lemma 1.10 that the topological closures¹⁰ $\overline{\Delta_f(\tau)}$ ¹¹ of the cones $\Delta_f(\tau)$ form a fan¹² $\overline{\Delta_f}$ of rational polyhedral cones¹³.

Remark 1.12. The function m_f from Definition 1.6 is linear on each cone $\overline{\Delta_f(\tau)}$.

Remark 1.13. Let \mathbf{f} and \mathcal{I} be as in Definition 1.2. Let $k \in \mathbf{R}_{\geq 0}^n$ and let τ and τ' be faces of $\Gamma_{\mathbf{f}}$ and $\Gamma_{\mathcal{I}}$, respectively. We then have analogous definitions and results for $m_{\mathbf{f}}(k)$, $F_{\mathbf{f}}(k)$, $\sim_{\mathbf{f}}$, $\Delta_{\mathbf{f}}(\tau)$, $\Delta_{\mathbf{f}}$ and $\overline{\Delta_{\mathbf{f}}}$, associated to \mathbf{f} , and for $m_{\mathcal{I}}(k)$, $F_{\mathcal{I}}(k)$, $\sim_{\mathcal{I}}$, $\Delta_{\mathcal{I}}(\tau')$, $\Delta_{\mathcal{I}}$ and $\overline{\Delta_{\mathcal{I}}}$, associated to \mathcal{I} .

We state without proofs, the following two lemmas (see e.g. [6]).

Lemma 1.14. *Let Δ be the cone strictly positively spanned by the vectors $k_1, \dots, k_r \in \mathbf{R}_{\geq 0}^n \setminus \{0\}$. Then there exists a finite partition of Δ into cones Δ_i , such that each Δ_i is strictly positively spanned by a \mathbf{R} -linearly independent subset of $\{k_1, \dots, k_r\}$. We call such a decomposition a simplicial decomposition of Δ without introducing new rays.*

Lemma 1.15. *Let Δ be a rational simplicial cone. Then there exists a finite partition of Δ into simple cones. (In general, such a decomposition requires the introduction of new rays.)*

Finally, we need the following notion, which is closely related to the notion of a simple cone.

⁵With respect to the standard topology of $\mathbf{R}_{\geq 0}^n$.

⁶A subset of $\mathbf{R}_{\geq 0}^n$ is called relatively open if it is open in its affine closure.

⁷A subset C of \mathbf{R}^n is called a convex cone if it is a convex set and $\lambda x \in C$ for all $x \in C$ and all $\lambda \in \mathbf{R}_{>0}$.

⁸A facet is a face of codimension one.

⁹The dimension of a convex cone is the dimension of its affine hull.

¹⁰With respect to the standard topology of $\mathbf{R}_{\geq 0}^n$.

¹¹ $\overline{\Delta_f(\tau)} = \{\lambda_1 k_1 + \lambda_2 k_2 + \dots + \lambda_r k_r \mid \lambda_i \in \mathbf{R}_{\geq 0}\} = \{k \in \mathbf{R}_{\geq 0}^n \mid F_f(k) \supset \tau\}$.

¹²A fan \mathcal{F} is a finite set of rational polyhedral cones such that every face of a cone in \mathcal{F} is contained in \mathcal{F} and the intersection of each two cones C and C' in \mathcal{F} is a face of both C and C' .

¹³A rational polyhedral cone is a closed convex cone, generated by a finite subset of \mathbf{Z}^n .

Definition 1.16. Let k_1, \dots, k_r be \mathbf{Q} -linearly independent vectors in \mathbf{Z}^n . The multiplicity of k_1, \dots, k_r , denoted by $\text{mult}(k_1, \dots, k_r)$, is defined as the index of the lattice $\mathbf{Z}k_1 + \dots + \mathbf{Z}k_r$ in the group of points with integral coordinates in the subspace spanned by k_1, \dots, k_r of the \mathbf{Q} -vector space \mathbf{Q}^n . One can check that this number equals the cardinality of the set

$$\mathbf{Z}^n \cap \left\{ \sum_{i=1}^r \lambda_i k_i \mid 0 \leq \lambda_i < 1 \text{ for } i = 1, \dots, r \right\}.$$

Let Δ be the cone strictly positively spanned by k_1, \dots, k_r . We define the multiplicity of Δ as the multiplicity of k_1, \dots, k_r , and we denote it by $\text{mult } \Delta$.

Remark 1.17. Let Δ be as in Definition 1.16. Note that Δ is simple if and only if $\text{mult } \Delta = \text{mult}(k_1, \dots, k_r) = 1$.

2. IGUSA'S LOCAL ZETA FUNCTION OF A MONOMIAL IDEAL AND A POLYNOMIAL MEASURE

In the previous section we met partitions Δ_f , $\Delta_{\mathbf{f}}$ and $\Delta_{\mathcal{I}}$ of $\mathbf{R}_{\geq 0}^n$, associated to a polynomial, a polynomial mapping and a monomial ideal, respectively. When, as in this section, we are dealing with two polynomials f and g , a polynomial mapping \mathbf{f} and a polynomial g or an ideal \mathcal{I} and a polynomial g , we will also consider a partition of $\mathbf{R}_{\geq 0}^n$ by finitely and strictly positively spanned rational cones, associated to both f and g , both \mathbf{f} and g or both \mathcal{I} and g .

Definition 2.1 ($\Delta_{f,g}$, $\Delta_{\mathbf{f},g}$, $\Delta_{\mathcal{I},g}$). Let f , \mathbf{f} , \mathcal{I} and g be as in Definition 1.2. The partition $\Delta_{f,g}$ of $\mathbf{R}_{\geq 0}^n$, associated to f and g , will consist of all the nonempty intersections of cones in Δ_f with cones in Δ_g , i.e.

$$\Delta_{f,g} = \{ \Delta_f(\tau) \cap \Delta_g(\tau') \mid \begin{array}{l} \tau \text{ is a face of } \Gamma_f, \tau' \text{ is a face of } \Gamma_g \text{ and } \Delta_f(\tau) \cap \Delta_g(\tau') \neq \emptyset \end{array} \}.$$

The set of cones $\Delta_{f,g}$ will then be the quotient of $\mathbf{R}_{\geq 0}^n$ by the equivalence relation

$$k \sim_{f,g} k' \quad \text{if and only if} \quad F_f(k) = F_f(k') \text{ and } F_g(k) = F_g(k').$$

The partitions $\Delta_{\mathbf{f},g}$, associated to \mathbf{f} and g , and $\Delta_{\mathcal{I},g}$, associated to \mathcal{I} and g , are defined in the same way.

Example 2.2. Consider the monomial ideal $\mathcal{I} = (x^5y, x^3y^2, x^2y^5) \triangleleft \mathbf{Z}_p[x, y]$ and the polynomial $g(x, y) = x^4y^2 + xy^5 \in \mathbf{Z}_p[x, y]$. The Newton polyhedra $\Gamma_{\mathcal{I}}$ and Γ_g and the partitions $\Delta_{\mathcal{I}}$, Δ_g and $\Delta_{\mathcal{I},g}$ of $\mathbf{R}_{\geq 0}^2$ are drawn in Figure 1. Note that the vectors v_1, \dots, v_4 , v'_1, \dots, v'_3 , w_1, \dots, w_5 are perpendicular to the faces associated to the rays¹⁴ they span. One checks that g is non-degenerated over \mathbf{F}_p with respect to its Newton polyhedron, if and only if $p \neq 3$.

Let \mathcal{I} be a nonzero proper monomial ideal of $\mathbf{Z}_p[x_1, \dots, x_n]$ and g a non-zero polynomial in $\mathbf{Z}_p[x_1, \dots, x_n]$ with $g(0) = 0$, which is non-degenerated over \mathbf{F}_p with respect to all the faces of its Newton polyhedron. We consider the Newton polyhedra $\Gamma_{\mathcal{I}}$ and Γ_g of \mathcal{I} and g and the corresponding partitions $\Delta_{\mathcal{I}}$, Δ_g and $\Delta_{\mathcal{I},g}$ of $\mathbf{R}_{\geq 0}^n$.

¹⁴A ray is a cone of dimension one.






(a) $\Gamma_{\mathcal{I}}$ and its faces(b) Partition $\Delta_{\mathcal{I}}$ of $\mathbf{R}_{\geq 0}^n$ (c) Γ_g and its faces(d) Partition Δ_g of $\mathbf{R}_{\geq 0}^n$ (e) Partition $\Delta_{\mathcal{I},g}$ of $\mathbf{R}_{\geq 0}^n$

FIGURE 1. Newton polyhedra and partitions of $\mathbf{R}_{\geq 0}^n$ associated to $\mathcal{I} = (x^5y, x^3y^2, x^2y^5)$ and $g = x^4y^2 + xy^5$

Notation 2.3. From the definition of $\Delta_{\mathcal{I},g}$ and the fact that $\Delta_{\mathcal{I}}$ and Δ_g consist of disjoint sets, it follows that each cone δ in $\Delta_{\mathcal{I},g}$ can be written in a unique way as the intersection $\delta = \Delta_{\mathcal{I}}(\tau) \cap \Delta_g(\tau')$ of a cone $\Delta_{\mathcal{I}}(\tau)$ in $\Delta_{\mathcal{I}}$ with a cone $\Delta_g(\tau')$ in Δ_g . So, to each cone $\delta \in \Delta_{\mathcal{I},g}$ we can associate a face τ of $\Gamma_{\mathcal{I}}$ and a face τ' of Γ_g . For $\delta \in \Delta_{\mathcal{I},g}$, denote by g_δ the polynomial¹⁵ $g_{\tau'}$, with τ' the face of Γ_g associated to δ . Finally, for $\delta \in \Delta_{\mathcal{I},g}$, put

$$N_\delta = \# \{a \in (\mathbf{F}_p^\times)^n \mid \overline{g_\delta}(a) = 0\}.$$

Notation 2.4. For $k = (k_1, \dots, k_n) \in \mathbf{R}^n$, we denote

$$\sigma(k) = \sum_{i=1}^n k_i.$$

Theorem 2.5. *Let \mathcal{I} and g be as above. Then we have:*

$$Z_{\mathcal{I},gdx}(s) = \sum_{\delta \in \Delta_{\mathcal{I},g}} L_\delta S_\delta,$$

with

$$L_\delta = p^{-n} \left((p-1)^n - N_\delta \frac{p}{p+1} \right)$$

and

$$S_\delta = \sum_{k \in \mathbf{N}^n \cap \delta} p^{-m_{\mathcal{I}}(k)s - m_g(k) - \sigma(k)},$$

for every cone δ in $\Delta_{\mathcal{I},g}$.

The above formula for S_δ can be rewritten as a rational expression in p^{-s} . Consider therefore a partition of the rational cone δ into rational simplicial cones δ_i , $i \in I$, without the introduction of new rays. For each i , let δ_i be the cone strictly positively spanned by the linearly independent, primitive vectors $k_{i,1}, \dots, k_{i,r_i} \in \mathbf{N}^n \setminus \{0\}$. Then

$$S_\delta = \sum_{i \in I} \frac{\sum_h p^{m_{\mathcal{I}}(h)s + m_g(h) + \sigma(h)}}{\prod_{j=1}^{r_i} (p^{m_{\mathcal{I}}(k_{i,j})s + m_g(k_{i,j}) + \sigma(k_{i,j})} - 1)},$$

where h runs through the elements of the set

$$\mathbf{Z}^n \cap \left\{ \sum_{j=1}^{r_i} \lambda_j k_{i,j} \mid 0 \leq \lambda_j < 1 \text{ for } j = 1, \dots, r_i \right\}.$$

From the formula for L_δ and the rational expression for S_δ , it then follows that the real parts of the candidate-poles of $Z_{\mathcal{I},gdx}$ are given by the rational numbers

$$-\frac{m_g(k) + \sigma(k)}{m_{\mathcal{I}}(k)},$$

for k a primitive generator of a ray in $\Delta_{\mathcal{I},g}$.

Proof. The proof is similar to the proofs of [6, Theorem 4.2] and [9, Proposition 2.1]. First we divide the integration domain \mathbf{Z}_p^n , based on the partition $\Delta_{\mathcal{I},g}$ of $\mathbf{R}_{\geq 0}^n$ associated to \mathcal{I} and g :

$$Z_{\mathcal{I},gdx}(s) = \int_{\mathbf{Z}_p^n} |\mathcal{I}(x)|^s |g(x)| |dx|$$

¹⁵See Definition 1.4.

primitive generator k of a ray in $\Delta_{\mathcal{I},g}$	$m_{\mathcal{I}}(k)$	$m_g(k)$	$\sigma(k)$	candidate-pole associated to k
(1, 0)	2	1	1	-1
(3, 1)	11	8	4	-12/11
(1, 1)	5	6	2	-8/5
(1, 2)	7	8	3	-11/7
(0, 1)	1	2	1	-3
h	$m_{\mathcal{I}}(h)$	$m_g(h)$	$\sigma(h)$	
(2, 1)	8	7	3	

TABLE 1. Data associated to (2, 1) and the primitive generators of the rays in $\Delta_{\mathcal{I},g}$.

$$= \sum_{\delta \in \Delta_{\mathcal{I},g}} \sum_{k \in \mathbb{N}^n \cap \delta} \int_{\substack{x \in \mathbf{Z}_p^n \\ \text{ord } x = k}} |\mathcal{I}(x)|^s |g(x)| |dx|.$$

Recall that $|\mathcal{I}(x)| = \max\{|f(x)| \mid f \in \mathcal{I}\} = p^{-\min\{\text{ord } f(x) \mid f \in \mathcal{I}\}}$. It's easy to see that this minimum is always attained in one of the monic monomials that generate \mathcal{I} . If x^ω is a monomial in $\mathbf{Z}_p[x_1, \dots, x_n]$, then for $x \in \mathbf{Z}_p^n$ with $\text{ord } x = k$, the order of x^ω equals $k \cdot \omega$. So, since $\min\{k \cdot \omega \mid \omega \in \text{supp}(\mathcal{I})\} = m_{\mathcal{I}}(k)$, it follows that

$$Z_{\mathcal{I},gdx}(s) = \sum_{\delta \in \Delta_{\mathcal{I},g}} \sum_{k \in \mathbb{N}^n \cap \delta} p^{-m_{\mathcal{I}}(k)s} \int_{\substack{x \in \mathbf{Z}_p^n \\ \text{ord } x = k}} |g(x)| |dx|.$$

From the proof of [6, Theorem 4.2], it follows that the integral in the previous equation is equal to $p^{-m_g(k)-\sigma(k)} L_\delta$. Hence,

$$Z_{\mathcal{I},gdx}(s) = \sum_{\delta \in \Delta_{\mathcal{I},g}} \sum_{k \in \mathbb{N}^n \cap \delta} p^{-m_{\mathcal{I}}(k)s} p^{-m_g(k)-\sigma(k)} L_\delta$$

and since L_δ is independent of k , we find

$$\begin{aligned} Z_{\mathcal{I},gdx}(s) &= \sum_{\delta \in \Delta_{\mathcal{I},g}} L_\delta \sum_{k \in \mathbb{N}^n \cap \delta} p^{-m_{\mathcal{I}}(k)s - m_g(k) - \sigma(k)} \\ &= \sum_{\delta \in \Delta_{\mathcal{I},g}} L_\delta S_\delta. \end{aligned}$$

For the easy proof of the rational formula for S_δ , we refer to [6, Theorem 4.2]. Essential here is the fact that the functions $m_{\mathcal{I}}$ and m_g are linear on each δ . ■

Example 2.6 (Continuation of Example 2.2). We will now calculate Igusa's local zeta function $Z_{\mathcal{I},gdx}$ for the monomial ideal \mathcal{I} and the polynomial g from Example 2.2. One can verify that g is non-degenerated over \mathbf{F}_p with respect to Γ_g , if and only if $p \neq 3$. So, we restrict to this case and use the formula from Theorem 2.5 to calculate $Z_{\mathcal{I},gdx}(s)$.

Tables 1 and 2 show the data related to the cones in $\Delta_{\mathcal{I},g}$ and the primitive generators of its rays, needed to 'fill in' our formula. The number N_g , appearing

cone δ	dim δ	primitive generators	mult δ	N_δ	L_δ	S_δ
δ_0	0			N_g	$\left(\frac{p-1}{p}\right)^2 - \frac{N_g}{p(p+1)}$	1
δ_1	1	(1, 0)		0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{p^{2s+2}-1}$
δ_2	2	(1, 0), (3, 1)	1	0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{(p^{2s+2}-1)(p^{11s+12}-1)}$
δ_3	1	(3, 1)		0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{p^{11s+12}-1}$
δ_4	2	(3, 1), (1, 1)	2	0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1+p^{8s+10}}{(p^{11s+12}-1)(p^{5s+8}-1)}$
δ_5	1	(1, 1)		N_g	$\left(\frac{p-1}{p}\right)^2 - \frac{N_g}{p(p+1)}$	$\frac{1}{p^{5s+8}-1}$
δ_6	2	(1, 1), (1, 2)	1	0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{(p^{5s+8}-1)(p^{7s+11}-1)}$
δ_7	1	(1, 2)		0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{p^{7s+11}-1}$
δ_8	2	(1, 2), (0, 1)	1	0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{(p^{7s+11}-1)(p^{s+3}-1)}$
δ_9	1	(0, 1)		0	$\left(\frac{p-1}{p}\right)^2$	$\frac{1}{p^{s+3}-1}$

TABLE 2. Data associated to the cones of $\Delta_{\mathcal{I},g}$.

in Table 2, is the number of elements in the locus $\{(x, y) \in (\mathbf{F}_p^\times)^2 \mid \overline{g}(x, y) = x^4y^2 + xy^5 = 0\}$. One can check that this number equals

$$N_g = \begin{cases} 3(p-1) & \text{if } p \equiv 1, 7 \pmod{12}, \\ p-1 & \text{if } p \equiv 2, 5, 11 \pmod{12}. \end{cases}$$

The calculations are relatively easy, because, since we work in dimension 2, all cones $\delta \in \Delta_{\mathcal{I},g}$ are simplicial. Moreover, all cones are simple, except δ_4 . Consequently, we have to take the single point

$$(2, 1) \in \mathbf{Z}^n \cap \{\lambda(3, 1) + \mu(1, 1) \mid 0 \leq \lambda, \mu < 1\}$$

into account, in the calculation of S_{δ_4} .

For $p \equiv 1, 7 \pmod{12}$, we find

$$Z_{\mathcal{I},gdx}(s) = \sum_{\delta \in \Delta_{\mathcal{I},g}} L_\delta S_\delta = \frac{p^6(p-1)A(p, t)}{(p+1)(p^2-t^2)(p^{12}-t^{11})(p^8-t^5)(p^{11}-t^7)(p^3-t)},$$

where $t = p^{-s}$ and $A(p, t)$ is the following polynomial in p and t .

$$\begin{aligned} A(p, t) = & (-p^2 - 3p + 1)t^{21} + (p^5 + 3p^4 - p^3)t^{20} + (-p^5 + p^4 + 4p^3 - p^2)t^{19} \\ & + (-p^7 - 3p^6 + p^5 - p^4 + p^2)t^{18} + (2p^7 - 2p^5)t^{17} + (p^6 - p^4)t^{16} \\ & + (-p^9 + p^7 - p^6 + p^4)t^{15} + (p^{13} + 3p^{12} - p^{11} + p^9 - p^7)t^{14} - 3p^{15}t^{13} \end{aligned}$$

$$\begin{aligned}
& + (-p^{15} - 3p^{14} + p^{13})t^{12} + (3p^{17} + p^{15} - p^{13})t^{11} + (-p^{18} + p^{16} + p^{14} + 3p^{13} - p^{12})t^{10} \\
& + (-2p^{17} - 3p^{16} + 2p^{15})t^9 + (p^{20} - p^{18} + 2p^{17} - 5p^{15})t^8 + (-p^{20} + 4p^{18})t^7 \\
& + (-p^{19} + p^{17})t^6 + (-p^{25} - 3p^{24} + p^{23})t^3 + 3p^{27}t^2 + 3p^{26}t + p^{30} - 3p^{29} - p^{28}
\end{aligned}$$

Note that none of the candidate-poles cancel in this case. For $p \equiv 2, 5, 11 \pmod{12}$, we find a similar result.

3. COMPUTATION OF AN IMPORTANT INTEGRAL USING HENSEL'S LEMMA

In order to prove our second main theorem in the next section, we will need a formula for the integral

$$\int_{(\mathbf{Z}_p^\times)^n} |f(x)|^s |g(x)| |dx|,$$

where f and g are polynomials over \mathbf{Z}_p in n variables. In Corollary 3.6 we will find a reasonable formula under a number of non-degeneracy conditions on f and g that will be explained further on in this section. The result demands some preparation involving Hensel's Lemma.

Lemma 3.1 (Hensel's lemma in several variables). *Let $f = (f_1, \dots, f_n)$ be an n -tuple of polynomials f_i in $\mathbf{Z}_p[x_1, \dots, x_n]$. Denote by $J(f) = |\partial f_i / \partial x_j|_{i,j}$ the Jacobian determinant of f . Let $k \in \mathbf{N} \setminus \{0\}$ and let $a = (a_1, \dots, a_n) \in \mathbf{Z}_p^n$ be a k -approximate root of f (i.e. $f_i(a) \equiv 0 \pmod{p^k}$ for all i) and suppose that $J(f, a) \not\equiv 0 \pmod{p}$. Then there exists a unique root $a' \in \mathbf{Z}_p^n$ of f near a (i.e. $f_i(a') = 0$ and $a'_i \equiv a_i \pmod{p^k}$ for all i).*

Proof. For the case $k = 1$, we refer to [4, Section 4.6, Theorem 2]. The result for $k > 1$ can be proved by applying the statement for $k = 1$ to the polynomial $g(y) = p^{-(k-1)} f(a + p^{k-1}y)$. ■

Corollary 3.2. *Cfr. [6, Corollary 3.4]. Let $f(x) \in \mathbf{Z}_p[x]$ and $k, l \in \mathbf{N} \setminus \{0\}$ with $k \geq l$. Let $a \in \mathbf{Z}_p$ such that $f(a) \equiv 0 \pmod{p^l}$ and $f'(a) \not\equiv 0 \pmod{p}$. Then there exists an element $a' \in \mathbf{Z}_p$ such that*

$$\{x \in a + p^l \mathbf{Z}_p \mid f(x) \equiv 0 \pmod{p^k}\} = a' + p^k \mathbf{Z}_p.$$

Proof. By the previous lemma, it follows that there exists a unique root $a' \in \mathbf{Z}_p$ of f which is congruent to a modulo p^l . One can verify that a' satisfies the required condition. ■

Corollary 3.3. *Let $f, g \in \mathbf{Z}_p[x_1, x_2]$ and $k \in \mathbf{N} \setminus \{0\}$. Let $a = (a_1, a_2) \in \mathbf{Z}_p^2$ such that $f(a) \equiv g(a) \equiv 0 \pmod{p}$ and $|J(f, g, a)| \not\equiv 0 \pmod{p}$. Then there exists an element $a' = (a'_1, a'_2) \in \mathbf{Z}_p^2$ such that*

$$A_{k,k} = \{x = (x_1, x_2) \in a + (p\mathbf{Z}_p)^2 \mid f(x) \equiv g(x) \equiv 0 \pmod{p^k}\} = a' + (p^k \mathbf{Z}_p)^2.$$

Proof. This follows in the same way as Corollary 3.2 from Lemma 3.1. ■

It follows that the set $A_{k,k} \subset \mathbf{Z}_p^2$ has measure p^{-2k} . We will need the following generalization of this result.

Lemma 3.4. *Let $n \geq 2$ and $f, g \in \mathbf{Z}_p[x_1, \dots, x_n]$. Let $k, l \in \mathbf{N} \setminus \{0\}$ with $k \geq l$. Let $a = (a_1, \dots, a_n) \in \mathbf{Z}_p^n$ such that $f(a) \equiv g(a) \equiv 0 \pmod{p}$ and such that the Jacobian matrix $J(f, g, a)$ has rank 2 modulo p . Then the measure of the set*

$$A_{k,l} = \{x = (x_1, \dots, x_n) \in a + (p\mathbf{Z}_p)^n \mid f(x) \equiv 0 \pmod{p^k} \text{ and } g(x) \equiv 0 \pmod{p^l}\}$$

equals $p^{-n-k-l+2}$.

Proof. Since $J(f, g, a)$ has rank 2 modulo p , we can assume, without loss of generality, that

$$\begin{vmatrix} \frac{\partial f}{\partial x_1}(a) & \frac{\partial f}{\partial x_2}(a) \\ \frac{\partial g}{\partial x_1}(a) & \frac{\partial g}{\partial x_2}(a) \end{vmatrix} \not\equiv 0 \pmod{p}.$$

This also implies that $(\partial f/\partial x_1)(a)$ and $(\partial f/\partial x_2)(a)$ are not both congruent to zero modulo p . Let us again assume that $(\partial f/\partial x_1)(a) \not\equiv 0 \pmod{p}$.

Fix $a'_3, \dots, a'_n \in \mathbf{Z}_p$ with $a'_i \equiv a_i \pmod{p}$ for $i = 3, \dots, n$. Applying Corollary 3.3 to the polynomials $f(x_1, x_2, a'_3, \dots, a'_n)$ and $g(x_1, x_2, a'_3, \dots, a'_n)$ in the variables x_1, x_2 and to $(a_1, a_2) \in \mathbf{Z}_p^2$, we find that there exist $a'_1, a'_2 \in \mathbf{Z}_p$, depending on a'_3, \dots, a'_n , such that for all $x_3, \dots, x_n \in \mathbf{Z}_p$, satisfying $x_i \equiv a'_i \pmod{p^k}$ for $i = 3, \dots, n$, one has

$$\begin{aligned} \{(x_1, x_2) \in (a_1, a_2) + (p\mathbf{Z}_p)^2 \mid f(x_1, \dots, x_n) \equiv g(x_1, \dots, x_n) \equiv 0 \pmod{p^l}\} \\ = (a'_1, a'_2) + (p^l\mathbf{Z}_p)^2. \end{aligned}$$

Consider now $a''_2 \in \mathbf{Z}_p$ with $a''_2 \equiv a'_2 \pmod{p^l}$. Applying Corollary 3.2 to the polynomial $f(x_1, a''_2, a'_3, \dots, a'_n) \in \mathbf{Z}_p[x_1]$ and $a'_1 \in \mathbf{Z}_p$, we find an element $a'_1 \in \mathbf{Z}_p$, depending on a''_2, a'_3, \dots, a'_n , such that for all $x_2, \dots, x_n \in \mathbf{Z}_p$, satisfying $x_2 \equiv a''_2 \pmod{p^l}$, $x_i \equiv a'_i \pmod{p^k}$ for $i = 3, \dots, n$, we have

$$\{x_1 \in a'_1 + p^l\mathbf{Z}_p \mid f(x_1, \dots, x_n) \equiv 0 \pmod{p^k}\} = a'_1 + p^k\mathbf{Z}_p.$$

These two considerations show that the set

$$A_{k,l} = \{x = (x_1, \dots, x_n) \in a + (p\mathbf{Z}_p)^n \mid f(x) \equiv 0 \pmod{p^k} \text{ and } g(x) \equiv 0 \pmod{p^l}\}$$

equals the union

$$(2) \quad \bigcup_{\substack{a'_3 + p^k\mathbf{Z}_p, \dots, a'_n + p^k\mathbf{Z}_p \\ a'_i \equiv a_i \pmod{p}}} \bigcup_{\substack{a''_2 + p^k\mathbf{Z}_p \\ a''_2 \equiv a'_2 \pmod{p^l}}} (a'_1, a''_2, a'_3, \dots, a'_n) + (p^k\mathbf{Z}_p)^n,$$

where the first union is taken over all elements $(a'_3 + p^k\mathbf{Z}_p, \dots, a'_n + p^k\mathbf{Z}_p) \in (\mathbf{Z}_p/p^k\mathbf{Z}_p)^{n-2}$ with $a'_i \equiv a_i \pmod{p}$ for $i = 3, \dots, n$ and the second union is over all the cosets $a''_2 + p^k\mathbf{Z}_p \in \mathbf{Z}_p/p^k\mathbf{Z}_p$ with $a''_2 \equiv a'_2 \pmod{p^l}$, where a'_2 depends on a'_3, \dots, a'_n and a'_1 depends on a''_2, a'_3, \dots, a'_n as described above.

Because the union (2) is a disjoint union, we now find easily that the measure of $A_{k,l}$ equals $p^{(k-1)(n-2)}p^{k-l}p^{-kn} = p^{-n-k-l+2}$. \blacksquare

Now we can prove the following proposition, which is the key result for the formula we are looking for.

Proposition 3.5. *Let $n \geq 2$ and $f, g \in \mathbf{Z}_p[x_1, \dots, x_n]$. Let $a \in \mathbf{Z}_p^n$ be no solution to any of the following sets of congruence relations:*

$$(3) \quad \begin{cases} f(x) \equiv 0 \pmod{p} \\ \frac{\partial f}{\partial x_i}(x) \equiv 0 \pmod{p}, \quad i = 1, \dots, n, \end{cases} \quad \begin{cases} g(x) \equiv 0 \pmod{p} \\ \frac{\partial g}{\partial x_i}(x) \equiv 0 \pmod{p}, \quad i = 1, \dots, n \end{cases}$$

and $\begin{cases} f(x) \equiv 0 \pmod{p} \\ g(x) \equiv 0 \pmod{p} \\ \left| \begin{array}{cc} \frac{\partial f}{\partial x_i}(a) & \frac{\partial f}{\partial x_j}(a) \\ \frac{\partial g}{\partial x_i}(a) & \frac{\partial g}{\partial x_j}(a) \end{array} \right| \equiv 0 \pmod{p}, \quad i, j = 1, \dots, n; \ i < j. \end{cases}$

Then for $s \in \mathbf{C}$ with $\Re(s) > 0$, we have

$$\int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| = \begin{cases} p^{-n} & \text{if } f(a) \not\equiv 0 \text{ and } g(a) \not\equiv 0 \pmod{p}, \\ p^{-n} \frac{p-1}{p^{s+1}-1} & \text{if } f(a) \equiv 0 \text{ and } g(a) \not\equiv 0 \pmod{p}, \\ p^{-n} \frac{1}{p+1} & \text{if } f(a) \not\equiv 0 \text{ and } g(a) \equiv 0 \pmod{p}, \\ p^{-n} \frac{p-1}{(p^{s+1}-1)(p+1)} & \text{if } f(a) \equiv g(a) \equiv 0 \pmod{p}. \end{cases}$$

Proof. The first three cases follow easily from [6, Proposition 3.1]. Let us now assume $f(a) \equiv g(a) \equiv 0 \pmod{p}$. For $x \in a + (p\mathbf{Z}_p)^n$, we have $f(x) \equiv f(a) \equiv g(x) \equiv g(a) \equiv 0 \pmod{p}$. The order of both $f(x)$ and $g(x)$ is thus at least one and we can write

$$\begin{aligned} & \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| \\ &= \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \int_{\substack{x \in a+(p\mathbf{Z}_p)^2 \\ \text{ord } f(x)=k, \text{ ord } g(x)=l}} |f(x)|^s |g(x)| |dx| \\ &= \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} p^{-ks} p^{-l} \cdot \mu(\{x \in a + (p\mathbf{Z}_p)^2 \mid \text{ord } f(x) = k \text{ and } \text{ord } g(x) = l\}). \end{aligned}$$

We will prove below that the measure of

$$(4) \quad \{x \in a + (p\mathbf{Z}_p)^2 \mid \text{ord } f(x) = k \text{ and } \text{ord } g(x) = l\}$$

equals $p^{-n-k-l}(p-1)^2$. Assuming it now, we can continue:

$$\begin{aligned} \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| &= \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} p^{-ks-l} p^{-n-k-l} (p-1)^2 \\ &= p^{-n} (p-1)^2 \sum_{k=1}^{\infty} p^{-(s+1)k} \sum_{l=1}^{\infty} p^{-2l} \\ &= p^{-n} (p-1)^2 \frac{p^{-(s+1)}}{1-p^{-(s+1)}} \frac{p^{-2}}{1-p^{-2}} \\ &= p^{-n} \frac{p-1}{(p^{s+1}-1)(p+1)}, \end{aligned}$$

giving the result.

It remains to prove that the measure of (4) is equal to $p^{-n-k-l}(p-1)^2$. With the notation of Lemma 3.4, we have that $A_{k,l} \supset A_{k+1,l}$ and $A_{k,l} \supset A_{k,l+1}$ and that the set $\{x \in a + (p\mathbf{Z}_p)^2 \mid \text{ord } f(x) = k \text{ and } \text{ord } g(x) = l\}$ is the complement of

$A_{k+1,l} \cup A_{k,l+1}$ in $A_{k,l}$. Noting further that $A_{k+1,l} \cap A_{k,l+1} = A_{k+1,l+1}$ and making use of Lemma 3.4, we can calculate that

$$\begin{aligned} & \mu(\{x \in a + (p\mathbf{Z}_p)^2 \mid \text{ord } f(x) = k \text{ and } \text{ord } g(x) = l\}) \\ &= \mu(A_{k,l}) - \mu(A_{k+1,l}) - \mu(A_{k,l+1}) + \mu(A_{k+1,l+1}) \\ &= p^{-n-k-l+2} - 2p^{-n-k-l+1} + p^{-n-k-l} \\ &= p^{-n-k-l}(p-1)^2. \end{aligned}$$

■

Corollary 3.6. *Let $n \geq 2$ and $f, g \in \mathbf{Z}_p[x_1, \dots, x_n]$. Put*

$$\begin{aligned} N &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \bar{f}(a) = 0 \text{ and } \bar{g}(a) \neq 0\} \\ P &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \bar{f}(a) \neq 0 \text{ and } \bar{g}(a) = 0\} \\ Q &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \bar{f}(a) = \bar{g}(a) = 0\}. \end{aligned}$$

Suppose that none of the sets of congruence relations (3) from the previous proposition has a solution in $(\mathbf{Z}_p^\times)^n$. Then for $s \in \mathbf{C}$ with $\Re(s) > 0$, we have

$$\begin{aligned} & \int_{(\mathbf{Z}_p^\times)^n} |f(x)|^s |g(x)| |dx| = \\ & p^{-n} \left((p-1)^n - pN \frac{p^s - 1}{p^{s+1} - 1} - P \frac{p}{p+1} - pQ \frac{p^s(p+1) - 2}{(p^{s+1} - 1)(p+1)} \right). \end{aligned}$$

Proof. Splitting up the integration domain as follows, we get

$$\begin{aligned} & \int_{(\mathbf{Z}_p^\times)^n} |f(x)|^s |g(x)| |dx| = \\ & \sum_{\substack{a \in \{1, \dots, p-1\}^n \\ f(a) \not\equiv 0 \pmod p \\ g(a) \not\equiv 0 \pmod p}} \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| + \sum_{\substack{a \in \{1, \dots, p-1\}^n \\ f(a) \equiv 0 \pmod p \\ g(a) \not\equiv 0 \pmod p}} \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| \\ & + \sum_{\substack{a \in \{1, \dots, p-1\}^n \\ f(a) \not\equiv 0 \pmod p \\ g(a) \equiv 0 \pmod p}} \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx| + \sum_{\substack{a \in \{1, \dots, p-1\}^n \\ f(a) \equiv 0 \pmod p \\ g(a) \equiv 0 \pmod p}} \int_{a+(p\mathbf{Z}_p)^n} |f(x)|^s |g(x)| |dx|. \end{aligned}$$

Applying Proposition 3.5, we find

$$\begin{aligned} & \int_{(\mathbf{Z}_p^\times)^n} |f(x)|^s |g(x)| |dx| \\ &= ((p-1)^n - N - P - Q)p^{-n} + Np^{-n} \frac{p-1}{p^{s+1} - 1} + Pp^{-n} \frac{1}{p+1} \\ & \quad + Qp^{-n} \frac{p-1}{(p^{s+1} - 1)(p+1)} \\ &= p^{-n} \left((p-1)^n - pN \frac{p^s - 1}{p^{s+1} - 1} - P \frac{p}{p+1} - pQ \frac{p^s(p+1) - 2}{(p^{s+1} - 1)(p+1)} \right). \end{aligned}$$

■

4. IGUSA'S LOCAL ZETA FUNCTION OF A SINGLE POLYNOMIAL AND A POLYNOMIAL MEASURE

Let f, g be non-zero polynomials in $\mathbf{Z}_p[x_1, \dots, x_n]$ without a constant term. Suppose f and g are both non-degenerated¹⁶ over \mathbf{F}_p with respect to all the faces of their Newton polyhedron. We will consider the Newton polyhedra Γ_f and Γ_g of f and g and their corresponding partitions¹⁷ Δ_f , Δ_g and $\Delta_{f,g}$ of $\mathbf{R}_{\geq 0}^n$.

Notation 4.1. As before, each cone δ in $\Delta_{f,g}$ can be written in a unique way as the intersection $\delta = \Delta_f(\tau) \cap \Delta_g(\tau')$ of a cone $\Delta_f(\tau)$ in Δ_f with a cone $\Delta_g(\tau')$ in Δ_g . So, to each cone $\delta \in \Delta_{f,g}$ we can associate a face τ of Γ_f and a face τ' of Γ_g . For $\delta \in \Delta_{f,g}$, denote by f_δ the polynomial¹⁸ f_τ , with τ the face of Γ_f associated to δ . The polynomial g_δ is defined in the same way. Furthermore, for $\delta \in \Delta_{f,g}$, we put

$$N_\delta = \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{f_\delta}(a) = 0 \text{ and } \overline{g_\delta}(a) \neq 0\}$$

$$P_\delta = \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{f_\delta}(a) \neq 0 \text{ and } \overline{g_\delta}(a) = 0\}$$

$$Q_\delta = \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{f_\delta}(a) = \overline{g_\delta}(a) = 0\}.$$

Definition 4.2 (Non-degenerated). Let $n \geq 2$ and f, g be non-zero polynomials in $\mathbf{Z}_p[x_1, \dots, x_n]$ without a constant term. We say that the pair (f, g) is non-degenerated over \mathbf{F}_p with respect to all the cones in the partition $\Delta_{f,g}$ of $\mathbf{R}_{\geq 0}^n$, associated to f and g , if for every cone $\delta \in \Delta_{f,g}$ and all $a \in (\mathbf{Z}_p^\times)^n$, such that $f_\delta(a) \equiv g_\delta(a) \equiv 0 \pmod{p}$, the Jacobian matrix $J(f_\delta, g_\delta, a)$ has rank 2 modulo p . This amounts to saying that for every cone $\delta \in \Delta_{f,g}$, the set of congruences

$$\left\{ \begin{array}{l} f_\delta(x) \equiv 0 \pmod{p} \\ g_\delta(x) \equiv 0 \pmod{p} \\ \left| \begin{array}{cc} \frac{\partial f_\delta}{\partial x_i}(a) & \frac{\partial f_\delta}{\partial x_j}(a) \\ \frac{\partial g_\delta}{\partial x_i}(a) & \frac{\partial g_\delta}{\partial x_j}(a) \end{array} \right| \equiv 0 \pmod{p}, \quad i, j = 1, \dots, n; \ i < j \end{array} \right.$$

has no solutions in $(\mathbf{Z}_p^\times)^n$.

Theorem 4.3. Let p be a prime number and $n \geq 2$. Let f, g be non-zero polynomials in $\mathbf{Z}_p[x_1, \dots, x_n]$ with $f(0) = 0$ and $g(0) = 0$. Suppose that f and g are both non-degenerated over \mathbf{F}_p with respect to all the faces of their Newton polyhedron and that (f, g) is non-degenerated over \mathbf{F}_p with respect to all the cones in the partition $\Delta_{f,g}$ of $\mathbf{R}_{\geq 0}^n$, associated to f and g . Then we have:

$$Z_{f,gdx}(s) = \sum_{\delta \in \Delta_{f,g}} L_\delta S_\delta,$$

with

$$L_\delta = p^{-n} \left((p-1)^n - pN_\delta \frac{p^s - 1}{p^{s+1} - 1} - P_\delta \frac{p}{p+1} - pQ_\delta \frac{p^s(p+1) - 2}{(p^{s+1} - 1)(p+1)} \right)$$

and

$$S_\delta = \sum_{k \in \mathbf{N}^n \cap \delta} p^{-m_f(k)s - m_g(k) - \sigma(k)},$$

for every cone δ in $\Delta_{f,g}$.

¹⁶See Definition 1.4.

¹⁷See Definitions 1.9 and 2.1.

¹⁸See Definition 1.4.

Just as in Theorem 2.5, the S_δ can be calculated by considering a partition of δ into rational simplicial cones δ_i , $i \in I$, without the introduction of new rays. If for each i , we take δ_i to be strictly positively spanned by the linearly independent, primitive vectors $k_{i,1}, \dots, k_{i,r_i} \in \mathbf{N}^n \setminus \{0\}$, then we have

$$S_\delta = \sum_{i \in I} \frac{\sum_h p^{m_f(h)s + m_g(h) + \sigma(h)}}{\prod_{j=1}^{r_i} (p^{m_f(k_{i,j})s + m_g(k_{i,j}) + \sigma(k_{i,j})} - 1)},$$

where h runs through the elements of the set

$$\mathbf{Z}^n \cap \left\{ \sum_{j=1}^{r_i} \lambda_j k_{i,j} \mid 0 \leq \lambda_j < 1 \text{ for } j = 1, \dots, r_i \right\}.$$

The real parts of the candidate-poles for $Z_{f,gdx}$ are therefore given by the rational numbers -1 and

$$-\frac{m_g(k) + \sigma(k)}{m_f(k)}$$

for k a primitive generator of a ray in $\Delta_{f,g}$.

Proof. The proof is again similar to the proof of [6, Theorem 4.2]. Based on the partition $\Delta_{f,g}$ of $\mathbf{R}_{\geq 0}^n$, we split up the integration domain as follows:

$$\begin{aligned} Z_{f,gdx}(s) &= \int_{\mathbf{Z}_p^n} |f(x)|^s |g(x)| |dx| \\ &= \sum_{\delta \in \Delta_{f,g}} \sum_{k \in \mathbf{N}^n \cap \delta} \int_{\substack{x \in \mathbf{Z}_p^n \\ \text{ord } x = k}} |f(x)|^s |g(x)| |dx|. \end{aligned}$$

To calculate the integral over $\{x \in \mathbf{Z}_p^n \mid \text{ord } x = k\}$, we will make a change of variables. For $\delta \in \Delta_{f,g}$, $k \in \mathbf{N}^n \cap \delta$ and $x \in \mathbf{Z}_p^n$ with $\text{ord } x = k$, put $x_j = p^{k_j} u_j$ with $u_j \in \mathbf{Z}_p^\times$. Then

$$\begin{aligned} |dx| &= p^{-\sigma(k)} |du| \quad \text{and} \\ x^\omega &= x_1^{\omega_1} \cdots x_n^{\omega_n} = p^{k \cdot \omega} u^\omega. \end{aligned}$$

Note that $f(x)$ is a \mathbf{Z}_p -linear combination of the x^ω with $\omega \in \text{supp}(f)$ and thus of the $p^{k \cdot \omega} u^\omega$ in the new variables. For k fixed and $\omega \in \text{supp}(f)$, the scalar product $k \cdot \omega$ is minimal with value $m_f(k)$ for $\omega \in F_f(k) = \tau$, where τ is the face of Γ_f associated to δ , and not minimal outside τ (Cfr. Definitions 1.6, 1.7 and 1.9). Separating the maximal power of p , we can write

$$f(x) = p^{m_f(k)} (f_\tau(u) + p \tilde{f}_{\tau,k}(u)) = p^{m_f(k)} (f_\delta(u) + p \tilde{f}_{\tau,k}(u)),$$

with $\tilde{f}_{\tau,k}(u)$ a polynomial in $\mathbf{Z}_p[u_1, \dots, u_n]$, depending on f , τ and k . Analogously, we can write $g(x)$ as

$$g(x) = p^{m_g(k)} (g_{\tau'}(u) + p \tilde{g}_{\tau',k}(u)) = p^{m_g(k)} (g_\delta(u) + p \tilde{g}_{\tau',k}(u)).$$

This all leads to $Z_{f,gdx}(s) =$

$$\sum_{\delta \in \Delta_{f,g}} \sum_{k \in \mathbf{N}^n \cap \delta} p^{-m_f(k)s - m_g(k) - \sigma(k)} \int_{(\mathbf{Z}_p^\times)^n} |f_\delta(u) + p \tilde{f}_{\tau,k}(u)|^s |g_\delta(u) + p \tilde{g}_{\tau',k}(u)| |du|.$$

Let us put

$$L_\delta = \int_{(\mathbf{Z}_p^\times)^n} |f_\delta(u) + p\tilde{f}_{\tau,k}(u)|^s |g_\delta(u) + p\tilde{g}_{\tau',k}(u)| |du|.$$

Because of the non-degeneracy conditions on f and g , for every $\delta \in \Delta_{f,g}$ the polynomials f_δ and g_δ satisfy the condition formulated in the statement of Corollary 3.6. It follows immediately that the polynomials $f_\delta(u) + p\tilde{f}_{\tau,k}(u)$ and $g_\delta(u) + p\tilde{g}_{\tau',k}(u)$ satisfy the same condition. So by Corollary 3.6, we know that

$$\begin{aligned} L_\delta &= \int_{(\mathbf{Z}_p^\times)^n} |f_\delta(u) + p\tilde{f}_{\tau,k}(u)|^s |g_\delta(u) + p\tilde{g}_{\tau',k}(u)| |du| \\ &= p^{-n} \left((p-1)^n - pN_\delta \frac{p^s - 1}{p^{s+1} - 1} - P_\delta \frac{p}{p+1} - pQ_\delta \frac{p^s(p+1) - 2}{(p^{s+1} - 1)(p+1)} \right). \end{aligned}$$

Because this expression for L_δ is independent of k , we can write

$$Z_{f,gdx}(s) = \sum_{\delta \in \Delta_{f,g}} L_\delta \sum_{k \in \mathbf{N}^n \cap \delta} p^{-m_f(k)s - m_g(k) - \sigma(k)} = \sum_{\delta \in \Delta_{f,g}} L_\delta S_\delta,$$

giving the main formula. The ‘closed’ formula for S_δ , based on simplicial subdivision of δ , can be found following the argumentation of [6, Theorem 4.2]. \blacksquare

5. IGUSA’S LOCAL ZETA FUNCTION OF A POLYNOMIAL MAPPING AND A POLYNOMIAL MEASURE

Let $f_1, \dots, f_t, g \in \mathbf{Z}_p[x_1, \dots, x_n]$, such that $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$ is a nonconstant polynomial mapping, satisfying $\mathbf{f}(0) = 0$. In this section, we give an explicit formula for Igusa’s local zeta function $Z_{\mathbf{f},gdx}$, associated to \mathbf{f} and the integration measure $|g(x)||dx|$ on \mathbf{Z}_p^n , in the same style as the previous ones. This time we adapt a formula for Igusa’s local zeta function of a polynomial mapping, given in [15], which is a generalization of the formulas proven in [6] and [9]. In the same way the formula we will state here, generalizes the main results of Sections 2 and 4. Again, we restrict to the case where \mathbf{f} and g satisfy a number of non-degeneracy conditions (see below).

Because the derivation of the formula is completely analogous to what we have done in Sections 3 and 4 for a single polynomial, in this section, we will only give the results and omit the proofs. We start by stating the analogues of some results from Section 3, to conclude with the formula itself in Theorem 5.6.

Lemma 5.1. *Let \mathbf{f} and g be as above and suppose $n \geq t+1$. Let $k, l \in \mathbf{N} \setminus \{0\}$ and $a = (a_1, \dots, a_n) \in \mathbf{Z}_p^n$, such that $\mathbf{f}(a) \equiv 0 \pmod{p}$ and $g(a) \equiv 0 \pmod{p}$, and such that the Jacobian matrix $J(\mathbf{f}, g, a) = J(f_1, \dots, f_t, g, a)$ has rank $t+1$ modulo p . Then the measure of the set*

$$A_{k,l} = \{x = (x_1, \dots, x_n) \in a + (p\mathbf{Z}_p)^n \mid \mathbf{f}(x) \equiv 0 \pmod{p^k} \text{ and } g(x) \equiv 0 \pmod{p^l}\}$$

equals $p^{-n-(k-1)t-l+1}$.

Proposition 5.2. *Let \mathbf{f} , g and n be as above. Let $a \in \mathbf{Z}_p^n$, such that the following three conditions hold: if a is in the locus of \mathbf{f} , g and (\mathbf{f}, g) , respectively, then the Jacobian matrix of \mathbf{f} , g , (\mathbf{f}, g) , respectively, and a , has maximal rank (\star) . Then for*

$s \in \mathbf{C}$ with $\Re(s) > 0$, we have

$$\int_{a+(p\mathbf{Z}_p)^n} \|\mathbf{f}(x)\|^s |g(x)| dx = \begin{cases} p^{-n} & \text{if } \mathbf{f}(a) \not\equiv 0 \text{ and } g(a) \not\equiv 0 \pmod{p}, \\ p^{-n} \frac{p^t-1}{p^{s+t}-1} & \text{if } \mathbf{f}(a) \equiv 0 \text{ and } g(a) \not\equiv 0 \pmod{p}, \\ p^{-n} \frac{1}{p+1} & \text{if } \mathbf{f}(a) \not\equiv 0 \text{ and } g(a) \equiv 0 \pmod{p}, \\ p^{-n} \frac{p^t-1}{(p^{s+t}-1)(p+1)} & \text{if } \mathbf{f}(a) \equiv 0 \text{ and } g(a) \equiv 0 \pmod{p}. \end{cases}$$

Corollary 5.3. *Let \mathbf{f} , g and n be as above. Put*

$$\begin{aligned} N &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}}(a) = 0 \text{ and } \overline{g}(a) \neq 0\} \\ P &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}}(a) \neq 0 \text{ and } \overline{g}(a) = 0\} \\ Q &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}}(a) = 0 \text{ and } \overline{g}(a) = 0\}. \end{aligned}$$

Suppose that the three conditions (\star) from the previous proposition hold for all $a \in (\mathbf{Z}_p^\times)^n$. Then for $s \in \mathbf{C}$ with $\Re(s) > 0$, we have

$$\int_{(\mathbf{Z}_p^\times)^n} \|\mathbf{f}(x)\|^s |g(x)| dx = p^{-n} \left((p-1)^n - p^t N \frac{p^s-1}{p^{s+t}-1} - P \frac{p}{p+1} - pQ \frac{p^{t-1}(p^s(p+1)-1)-1}{(p^{s+t}-1)(p+1)} \right).$$

Notation 5.4. Again each cone δ in $\Delta_{\mathbf{f},g}$ can be written in a unique way as the intersection $\delta = \Delta_{\mathbf{f}}(\tau) \cap \Delta_g(\tau')$ of a cone $\Delta_{\mathbf{f}}(\tau)$ in $\Delta_{\mathbf{f}}$ with a cone $\Delta_g(\tau')$ in Δ_g . So, to each cone $\delta \in \Delta_{\mathbf{f},g}$ we can associate a face τ of $\Gamma_{\mathbf{f}}$ and a face τ' of Γ_g . For $\delta \in \Delta_{\mathbf{f},g}$, denote by \mathbf{f}_δ the polynomial \mathbf{f}_τ , with τ the face of $\Gamma_{\mathbf{f}}$ associated to δ , and analogously for g_δ . Furthermore, for $\delta \in \Delta_{\mathbf{f},g}$, we put

$$\begin{aligned} N_\delta &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}_\delta}(a) = 0 \text{ and } \overline{g_\delta}(a) \neq 0\} \\ P_\delta &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}_\delta}(a) \neq 0 \text{ and } \overline{g_\delta}(a) = 0\} \\ Q_\delta &= \#\{a \in (\mathbf{F}_p^\times)^n \mid \overline{\mathbf{f}_\delta}(a) = 0 \text{ and } \overline{g_\delta}(a) = 0\}. \end{aligned}$$

We need the following notion of non-degeneracy.

Definition 5.5 (strongly non-degenerated). Let \mathbf{f} , g and n be as above. We say that the pair (\mathbf{f}, g) is strongly non-degenerated over \mathbf{F}_p with respect to (all the cones of) the partition $\Delta_{\mathbf{f},g}$ of $\mathbf{R}_{\geq 0}^n$, associated to \mathbf{f} and g , if for every cone $\delta \in \Delta_{\mathbf{f},g}$ and all $\overline{a} \in (\mathbf{F}_p^\times)^n$, such that $\overline{f_{1,\delta}}(\overline{a}) = \cdots = \overline{f_{t,\delta}}(\overline{a}) = \overline{g_\delta}(\overline{a}) = 0$, the Jacobian matrix

$$J(\overline{\mathbf{f}_\delta}, \overline{g_\delta}, \overline{a}) = \begin{pmatrix} (\partial \overline{f_{1,\delta}} / \partial x_1)(\overline{a}) & \cdots & (\partial \overline{f_{1,\delta}} / \partial x_n)(\overline{a}) \\ \vdots & \ddots & \vdots \\ (\partial \overline{f_{t,\delta}} / \partial x_1)(\overline{a}) & \cdots & (\partial \overline{f_{t,\delta}} / \partial x_n)(\overline{a}) \\ (\partial \overline{g_\delta} / \partial x_1)(\overline{a}) & \cdots & (\partial \overline{g_\delta} / \partial x_n)(\overline{a}) \end{pmatrix}$$

has maximal rank $(= t+1)$.

Theorem 5.6. *Let p be a prime number. Let $\mathbf{f} = (f_1, \dots, f_t) : \mathbf{Q}_p^n \rightarrow \mathbf{Q}_p^t$ be a non-constant polynomial mapping, with $f_1, \dots, f_t \in \mathbf{Z}_p[x_1, \dots, x_n]$, satisfying $f_i(0) = 0$ for all i , and suppose that $n \geq t+1$. Let $g, g(0) = 0$ be a nonconstant polynomial in $\mathbf{Z}_p[x_1, \dots, x_n]$ that is non-degenerated over \mathbf{F}_p with respect to its Newton polyhedron. Suppose that \mathbf{f} and (\mathbf{f}, g) are strongly non-degenerated over \mathbf{F}_p with respect*

to the Newton polyhedron $\Gamma_{\mathbf{f}}$ and the partition $\Delta_{\mathbf{f},g}$ of $\mathbf{R}_{\geq 0}^n$, respectively. Then we have:

$$Z_{\mathbf{f},gdx}(s) = \sum_{\delta \in \Delta_{\mathbf{f},g}} L_{\delta} S_{\delta},$$

with

$$L_{\delta} = p^{-n} \left((p-1)^n - p^t N_{\delta} \frac{p^s - 1}{p^{s+t} - 1} - P_{\delta} \frac{p}{p+1} - p Q_{\delta} \frac{p^{t-1}(p^s(p+1) - 1) - 1}{(p^{s+t} - 1)(p+1)} \right)$$

and

$$S_{\delta} = \sum_{k \in \mathbf{N}^n \cap \delta} p^{-m_{\mathbf{f}}(k)s - m_g(k) - \sigma(k)},$$

for every cone δ in $\Delta_{\mathbf{f},g}$.

Again¹⁹ there exists a closed, rational expression for the S_{δ} , showing that the real parts of the candidate-poles for $Z_{\mathbf{f},gdx}$ are given by the rational numbers -1 and

$$-\frac{m_g(k) + \sigma(k)}{m_{\mathbf{f}}(k)}$$

for k a primitive generator of a ray in $\Delta_{\mathbf{f},g}$.

Proof. The proof of the theorem is completely similar to the proof of Theorem 4.3. ■

REFERENCES

- [1] E. Artal Bartolo, P. Cassou-Noguès, I. Luengo, and A. Melle Hernández, *Monodromy conjecture for some surface singularities*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 4, 605–640. MR MR1981174 (2004e:32030)
- [2] ———, *Quasi-ordinary power series and their zeta functions*, Mem. Amer. Math. Soc. **178** (2005), no. 841, vi+85. MR MR2172403 (2007d:14005)
- [3] B. Bories, *Zeta functions and bernstein-sato polynomials for ideals in dimension two*, Preprint (2010) available at <http://arxiv.org/abs/1012.5437>.
- [4] N. Bourbaki, *Commutative algebra. Chapters 1–7*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1989, Translated from the French, Reprint of the 1972 edition. MR MR979760 (90a:13001)
- [5] J. Denef, *The rationality of the Poincaré series associated to the p -adic points on a variety*, Invent. Math. **77** (1984), no. 1, 1–23. MR MR751129 (86c:11043)
- [6] J. Denef and K. Hoornaert, *Newton polyhedra and Igusa’s local zeta function*, J. Number Theory **89** (2001), no. 1, 31–64. MR MR1838703 (2002g:11170)
- [7] J. Denef and F. Loeser, *Caractéristiques d’Euler-Poincaré, fonctions zêta locales et modifications analytiques*, J. Amer. Math. Soc. **5** (1992), no. 4, 705–720. MR MR1151541 (93g:11118)
- [8] ———, *Motivic Igusa zeta functions*, J. Algebraic Geom. **7** (1998), no. 3, 505–537. MR MR1618144 (99j:14021)
- [9] J. Howald, M. Mustață, and C. Yuen, *On Igusa zeta functions of monomial ideals*, Proc. Amer. Math. Soc. **135** (2007), no. 11, 3425–3433 (electronic). MR MR2336554 (2008j:11177)
- [10] J.-i. Igusa, *Complex powers and asymptotic expansions. I. Functions of certain types*, J. Reine Angew. Math. **268/269** (1974), 110–130, Collection of articles dedicated to Helmut Hasse on his seventy-fifth birthday, II. MR MR0347753 (50 #254)
- [11] A. Némethi and W. Veys, *Generalized monodromy conjecture in dimension two*, Preprint (2010).
- [12] R. T. Rockafellar, *Convex analysis*, Princeton University Press, Princeton, N.J., 1970.
- [13] L. Van Proeyen and W. Veys, *Poles of the topological zeta function associated to an ideal in dimension two*, Math. Z. **260** (2008), no. 3, 615–627. MR 2434472 (2009f:14025)
- [14] ———, *The monodromy conjecture for zeta functions associated to ideals in dimension two*, Ann. Inst. Fourier (Grenoble) **60** (2010), no. 4, 1347–1362. MR 2722244

¹⁹Cfr. Theorems 2.5 and 4.3.

- [15] W. Veys and W. A. Zúñiga-Galindo, *Zeta functions for analytic mappings, log-principalization of ideals, and Newton polyhedra*, Trans. Amer. Math. Soc. **360** (2008), no. 4, 2205–2227. MR MR2366980 (2008i:11140)

KATHOLIEKE UNIVERSITEIT LEUVEN, DEPARTMENT OF MATHEMATICS, CELESTIJNENLAAN 200B,
3001 LEUVEN, BELGIUM

E-mail address: `bart.bories@wis.kuleuven.be`